

This protocol is structured in accordance with the CONSORT – statement for Randomized Controlled Trials with the extensions for Non-Pharmacological Treatment Interventions, Cluster Randomized Trial and Pragmatic Trials <http://www.consort-statement.org/home/>. Protocol sections are shown below each CONSORT item.

Research Protocol

May 2013

Poul Lundgaard Bak, Carsten Obel, Jin Liang Zhu

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|--|-----------------------|
| Title and abstract | | | |
| 1a Identification as a randomized trial in the title. 1b Structured summary of trial design, methods, results, and conclusions (for specific guidance see CONSORT for abstracts). | In the abstract, description of the experimental treatment, comparator, care providers, centers, and blinding status. | Identification as a cluster randomized trial in the title. | |

Resilience in schools - a pragmatic cluster randomized trial of a Brief Intervention Program.

| Content | Page |
|--|-------------|
| Background | |
| • The Child mental Health Research program and the intervention program Robusthed.dk | 3 |
| • The School Trial | 6 |
| Trial design & Intervention | 8 |
| Participant eligibility | 9 |
| Outcome measures | 10 |
| Sample size calculation | 11 |
| Randomization | 12 |
| Statistical Methods | 13 |
| Time table | 14 |
| Ethical and legal considerations | 14 |
| Project Feasibility | 14 |
| Qualifications of the research Group | 14 |
| Publication | 15 |
| Scientific perspectives | 15 |
| Practical perspectives, knowledge dissemination | 15 |
| Protocol access | 16 |
| Funding | 16 |
| References | 17 |
| Appendix A, letter trial group A | 19 |
| Appendix A1, parent information | 20 |
| Appendix B, letter trial group B | 21 |
| Appendix C, CONSORT publication items | 22 |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|---|---|--|
| Introduction - Background and objectives | | | |
| 2a Scientific background and explanation of rationale. 2b Specific objectives or hypotheses. | | 2a Rationale for using a cluster design. 2b Whether objectives pertain to the cluster level, the individual participant level or both. | Describe the health or health service problem that the intervention is intended to address and other interventions that may commonly be aimed at this problem. |

Background

The Child Mental Health Research Program and the interventions program Robusthed.dk

The Child Mental Health Research Program has been started by Tryg Foundation grants in 2012 and 2013 with the purpose of establishing high quality research projects.

The goals of the research program are:

1. To develop and test health care models which can strengthen the clinical and educational support of vulnerable children, families and surrounding professionals – facing the serious welfare challenges of today’s society. This includes the integration of information about the child’s strengths and difficulties with tailored interventions.
2. To elucidate early life causal patterns and consequences for long term mental and somatic health and integrate this knowledge in the support and intervention models.

The research program is organized in cooperation between Aarhus University and Region Midt. A description of the organization and the six integrated Work Packages of the program (WP1-6) can be found at the research program website ([link](#)).

The current project concerns a coherent brief intervention program ([Work Package 4: Step care interventions](#)) which has now been developed and is ready for testing in controlled trials.


The intervention program is designed to respond to calls to exploit low cost & large scale intervention delivery models expressed by for instance Professor Allan Kadin (Kadin & Blasé 2011) and Professors Anthony Roth and Peter Fonagy (Roth and Fonagy 2006). The idea is to meet the socio-economic pressure on the public welfare sector concerning mental health which is currently a considerable challenge in many countries (Foresight Mental Capital and Wellbeing Project 2008). At present the principal suggested support model for mentally vulnerable individuals and families consist of advice and training on individual, family and group basis by trained professionals. These are, however, resource demanding models of delivery which society has a hard time to honor. The situation speaks for the development and testing of more resource efficient models to supplement these traditional models. Cochrane reviews suggest that self-directed media and internet based intervention programs may be as effective as traditional models of delivery (Montgomery, Bjornstad & Dennis 2006). In some respects contemporary and future internet based programs is expected to be superior to the traditional models of delivery (O’Connell 2009, Perkins et al 2009). However, much more rigorous research is needed in the field.

The intervention program in this research project is called Robusthed.dk (Robusthed is the Danish word for Resilience¹) because the program is designed to support resilience development in mentally vulnerable children, young people and families in cooperation between professionals and parents. The program is designed to meet different levels of difficulties (with or without diagnoses) and can also be used in general health promotion, conflict management and prevention of bullying.

Children and adolescents need to develop resilience in order to handle the challenges of life, small as well as big ones – especially when being in a vulnerable position. Self-control and Self-regulation are important parts of being resilient. Self-control in childhood is a strong predictor for health, wellbeing and social behavior later in life (Moffitt 2011). Self-control is specifically addressed in the intervention program.

The program is web based and includes documented knowledge available from cognitive- and neuroscience ('The Social Brain') as well as established experiences from parent training programs and social learning research in line with the NICE guidelines (NICE 2008, 2009). This is transformed into simple coherent presentations in daily language – equally understandable for a student, a parent and a highly educated professional. The program can be used directly for children down to the age 6-7 and for younger children by the reinforcement of good parenting inherent in the program. The program exploits the opportunities of contemporary information technology including animations, podcasts and smartphone application for cognitive training and social support.

From a theoretical point of view the program is structured in 3 dimensions – which are coupled to specific problem solving from a list of 12 important types of life problems (e.g. conflict & bullying, anxiety, sadness, sleep, acute crisis, pain, dependency, learning difficulties, etc.):



A detailed review of the scientific background of the program is found on the Robusthed.dk website (on the English sub-site "About Us" and on the website of the Child Health Mental Research Program [here](#)).

Results published in 2012 from a related uncontrolled precursor project in which central elements from the current program was included (involving about 4000 teachers and preschool teachers working with children and families and 3000 parents) has demonstrated wide applicability and a high feasibility (Lundgaard Bak 2012).

¹ The English version webaddress is myresilience.org

The program can be tested by logging in on www.Robusthed.dk (The program is in Danish and in English):

User name: rpres

Password: rpres

In the intervention program all adults around the child are offered exactly the same background knowledge and practical knowledge of how resilience training of children and adolescents can be integrated in daily life. The purpose is to support the development of common knowledge and understanding about child development and provide opportunities to train coping situations with the child both at home and daycare /school. **The knowledge and tools of the program is communicated to the children and adolescents by 'their own adults'**. The program is designed to fit into ordinary curricular and extracurricular activities and contacts between families and professionals on group and/or individual level in order to minimize the need for extra resource input. The program is introduced for the target groups in standardized short lectures and courses and participants are given a trial-specific log-in to the program website.

The trials in the research project are designed to follow the 'Recommendation on Criteria for Establishing Strong Evidence of Effectiveness' from The National Academies report: Preventing Mental, Emotional, and Behavioral Disorders among Young People: Progress and Possibilities (O'Connell 2009).

It is reasonable to assume that resilience is valuable for human beings in almost any context. Therefore it is relevant to investigate the eventual impact of the intervention program in a range of different contexts.

Thus the aim is to investigate - with societal relevant primary outcome indicators - if the brief intervention program Robusthed.dk can have a significant positive impact on the lives of selected target groups of vulnerable children, young people and families. The target groups have been selected on the basis of the following criteria:

- Relevance with respect to problem severity and/or scale in society
- Representation of different age groups
- Representation of different organizational contexts
- Exploiting synergies in the Child Mental Health Research Program (see www.iupgrowth.com).
- Exploiting the opportunities of using administrative (register) data as outcome indicators as recommended by the Coalition for Evidence Based Policy (2012): *How Low-Cost Randomized Controlled Trials Are Possible in Many Areas of Social Policy*.

3 large scaled trials are started in 2013-14 – covering children and adolescents in care (0-18 year), school projects (6-17 year) and youth education (15-25 year). This protocol concerns the Children in Care trial.

Table 1: Target groups in current and future research projects.

| Age: | Pregnancy | 0-5 year | 6-17 year | 18-25 year |
|--|-----------|----------------------------------|--|-----------------|
| DK Interventions in current trials | | | | |
| Robusthed.dk | | Children and adolescents in care | | |
| | | | School intervention | |
| | | | | Youth Education |
| DK interventions covered by future Danish applications | | | | |
| Robusthed.dk | Stress | Young mothers | | |
| | | | ADHD | |
| International interventions covered by international applications | | | | |
| Robusthed.dk | | | Greenland: Suicide prevention (the project starts Summer 2013) | |
| | | | EU-Harvard/USA mental disorder | |

The School Trial

In this trial there is focus on supporting the collaboration between parents and teachers and others who deal with children every day by giving all of them introduction and access to the Resilience Program. This strategy is based on five lines of evidence:

1. The strategy is in line with decades of research on whole school approaches in health promotion exemplified by the WHO Health Promotion Schools project involving more than 40 countries (WHO 2006) and recommendations from a series of NICE reports (Adi et al 2007a, Adi et al 2007b, Shucksmith et al 2007, NICE 2008, 2009, Wyn 2000).
2. Longitudinal social network analyses over 20-30 years from the Framingham study has documented that smoking as well as smoking cessation, development of obesity and the experience of happiness spreads dynamically in large social networks (Christakis & Fowler 2007 & 2008, Fowler & Christakis 2008). This supports that it is a reasonable strategy to seed relevant knowledge simultaneously into the whole adult network around a group of (vulnerable) children and young people. The school is an optimal arena for this kind of intervention.
3. Based on the current trend in Denmark to include a much greater proportion of vulnerable children into ordinary classes it seems highly reasonable to engage the whole class/school network into the intervention – certainly because one of the themes in the program is prevention of bullying which often has vulnerable children as victims.
4. Based on existing medical internet research evidence, combining a web-based intervention with a social field model of delivery seems to be the best way to maximize internet program adherence (Mohr et al 2011, Neil et al 2009).
5. The pilot studies indicate that when members of the school community tries out the program on individual and/or group level with success, such local success stories stimulate the interest of others in the network to use the program (Poul Lundgaard Bak 2012).

Robusthed.dk fits into ordinary curricular and extracurricular activities such as parents meetings on group and/or individual level. Thus a minimum of extra resource input is needed. A typical intervention course will be:

1. 1-3 hour standardized introduction to Robusthed.dk to the professionals on the school (leaders, teachers, etc.). Afterwards the teachers introduce Robusthed.dk to parents on ordinary parent meeting (1-1½ hour) besides implementing the program in the schedule of the class. Alternatively the introduction is given to all professionals and parents on the school at the same time. The introductory lecture is also available on the website.
The purpose of the introduction is to give a short overview of the relevance and core ideas of Robusthed.dk and to raise engagement and curiosity to go on and explore the 'Robusthed.dk universe' needed for program fidelity afterwards.
Parents and staff members are given log-in to the Robusthed.dk website.
2. Teachers and parents select and use Robusthed.dk knowledge and tools appropriate for specific children and adolescents (individually/groups) integrated in daily activities as a general mental health promoting activity or with a focus on specific problem solving. If they wish to run a more formal general Robusthed.dk course for the kids, they may use the course proposals on the website (on the subsite 'Vitamins of Knowledge').
3. Teachers and parents can exchange experiences and ideas on individual/group meetings as a user driven process. Process intensity is defined by local needs.
4. Informed users may explore the scientific background for Robusthed.dk in the Science Review on the website (link on the subsite 'About Us').

Trial design & intervention

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---|-----------------------|
| Methods – Trial Design | | | |
| 3a Description of trial design (such as parallel, factorial) including allocation ratio. 3b Important changes to methods after trial commencement (such as eligibility criteria), with reasons. | | 3a Definition of cluster and description of how the design features apply to the clusters. | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|--|--|--|
| Methods – Interventions | | | |
| 5 The interventions for each group with sufficient details to allow replication, including how and when they were actually administered. | 4 Precise details of both the experimental treatment and comparator. 4a Description of the different components of the interventions and, when applicable, descriptions of the procedure for tailoring the interventions to individual participants. 4b Details of how the interventions were standardized. 4c Details of how adherence of care providers with the protocol was assessed or enhanced. | 5 Whether interventions pertain to the cluster level, the individual participant level or both. | Describe extra resources added to (or resources removed from) usual settings in order to implement intervention. Indicate if efforts were made to standardize the intervention or if the intervention and its delivery were allowed to vary between participants, practitioners, or study sites. Describe the comparator in similar detail to the intervention. |

This is a pragmatic cluster randomized trial. No methods will be changed after trial commencement. A cluster is here defined as a school.

In Denmark there are 1.377 public municipality schools for children and adolescents (0-9(10)) grade. All the schools are considered eligible for the trial. From this pool groups of schools are randomly selected by a computer program (STATA) for the trial:

- A control group of 30 schools.
- Intervention group A, 30 schools. In this group of schools a mail/letter is send to the school leader offering a login to the Resilience Program with a recommendation to give it to all staff members and all parents (The letter is shown in Appendix A (in danish). Info-letter to the parents is also included (see Appendix A.1).
- Intervention group B, 30 schools. In this group of schools a mail/letter is send to the school leader offering a login to the Resilience Program with a recommendation to give it to all staff members and all parents. Moreover an intro-course is offered to the school as described above in the background section (The letter is shown in Appendix B (in Danish). Info-letter to the parents is also included (see Appendix A.1).
- Intervention group C, 30 schools. In this group of schools a mail/letter is send to the school leader and the chairman of the board offering a login to the Resilience Program with a recommendation to give it to all staff members and all parents. Moreover an intro-course is offered to the school as described above in the background section (The content of the letter is the same as in group B). Info-letter to the parents is also included (see Appendix A.1).

Outside the random trial we have an intervention group D (about 20 schools) which is a group of schools selected by municipality managers in 3 large municipalities in Denmark. They are selected based on an assessment of their needs (which in this case typically is associated with a low income population and area). The intervention offered to this group of schools is the same as in the intervention groups B and C. The reason for the inclusion of this group in the research project is to study the impact of different pragmatic implementation strategies in terms of school participation rate and the impact of different low cost delivery models.

The intervention pertains both to the cluster level and individual participant level.

Participant eligibility

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|--|--|--|
| Methods - Participants | | | |
| 4a Eligibility criteria for participants. 4b Settings and locations where the data were collected. | 3 When applicable, eligibility criteria for centers and those performing the interventions. | 4a Eligibility criteria for clusters. | 3 Eligibility criteria should be explicitly framed to show the degree to which they include typical participants and/or, where applicable, typical providers (eg, nurses), institutions (eg, hospitals), communities (or localities eg, towns) and settings of care (eg, different healthcare financing systems). |

All 1.377 public municipality schools in Denmark are included in the pool from which control and intervention groups of schools are randomly selected. Thus all are eligible and no one is excluded.

Outcome measures

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---|---|
| Methods - Outcomes | | | |
| <p>6a Completely defined pre-specified primary and secondary outcome measures, including how and when they were assessed.</p> <p>6b Any changes to trial outcomes after the trial commenced, with reasons.</p> | | <p>6a Whether outcome measures pertain to the cluster level, the individual participant level or both.</p> | <p>Explain why the chosen outcomes and, when relevant, the length of follow-up are considered important to those who will use the results of the trial.</p> |

Primary trial outcome will be academic results which are measured in the same way in all schools in the country. Academic results are an important prognostic welfare indicator in itself and it also correlates with self-reported wellbeing and behavior (Ottosen et al 2010).

All Danish students are regularly tested on academic results in dynamic tests (adapted to the level of the student). The test is scored 1-100. The data are automatically collected in a central register in UNI-C which is a department in the Danish Ministry of Education (read more: www.uvm.dk).

We will use the scores from the National Tests in Danish reading in 2, 4, 6, and 8 grades as data-sets because Danish reading is a very fundamental competency for learning other disciplines as well and for social functioning in contemporary society. Moreover it is known that results in Danish reading are highly correlated to academic results in other disciplines. The change in score from tests before intervention to test scores after the intervention will be used as outcome variable. Data-sets in the years before (back to 2010) and after the intervention (2015 and 2016) will be included in the analyses. Based on the Recommendation on Criteria for Establishing Strong Evidence of Effectiveness (O’Connell 2009) follow up time will thus be at least 2-3 years - depending on available resources. The register data are expected to be nearly complete.

Intention to treat analysis will be the basic principle (not in intervention group D), provided the school participation rate is sufficiently high. If participation rate is low, schools in the intervention groups which accept participation will be compared to matched control schools as an efficacy analysis.

School absenteeism will be used as secondary outcome measure in the random trial.

The outcome measure pertains both to the cluster level and individual participant level.

In the intervention group D (which is outside the random trial) – exploratory analyses of more detailed data from the municipality registers will be conducted – e.g. School absenteeism and self-reported health behavior and wellbeing. As far as possible basic measures as SDQ (which is widely used in Denmark and many other countries) will be used. Depending on local data resources more detailed subgroup analyses on children with special needs will be conducted. In the communities where ‘Skolesundhed.dk’ is implemented (a part of the Child Mental Health Research Program work Package 3: Interactive epidemiology ([Link](#))) these data will be included in the analyses. In all cases datasets before and after intervention are included.

Sample size calculation

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|--|---|---|
| Methods – Sample Size | | | |
| <p>7a How sample size was determined.</p> <p>7b When applicable, explanation of any interim analyses and stopping guidelines.</p> | When applicable, details of whether and how the clustering by care providers or centers was addressed. | <p>7a Method of calculation, number of clusters(s) (and whether equal or unequal cluster sizes are assumed), cluster size, a coefficient of intra-cluster correlation (ICC or k), and an indication of its uncertainty.</p> | If calculated using the smallest difference considered important by the target decision maker audience (the minimally important difference) then report where this difference was obtained. |

The mean score of Danish Reading among Danish school children is estimated 6.5 with standard deviation 2.6 (Data from the [Databank](#) in the Ministry of Education). From a societal perspective we estimate that a 10% increase in the mean score after the trial would be considered to be sufficiently interesting for decision makers to eventually implement the program on a larger scale. If we choose significance level of 0.05 and power of 80%, we need 252 children in both trial group and control group.




Figure: Distribution of national test scores for Danish reading among children in public schools

The public schools have an average of 438 (sd 263) children. If we assume intra-cluster correlation coefficient (ICC) of 0.02, we need 7 schools in each trial group. Assuming that 80% of the selected schools will response and participate, we need to include 9 schools in each trial group.

This present trial is a much larger scale trial - not only to ensure maximum statistical power but also to investigate the level of feasibility of large scale and low cost delivery models.

Randomization and blinding

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|---|--|---|
| Methods – Randomization – sequence generation | | | |
| 8a Method used to generate the random allocation sequence. 8b Type of randomization; details of any restriction (such as blocking and block size) | When applicable, how care providers were allocated to each trial group. | 8b Details of stratification or matching if used. | |
| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
| Methods – Randomization – Allocation concealment mechanism | | | |
| 9 Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned. | | Specification that allocation was based on clusters rather than individuals and whether allocation concealment (if any) was at the cluster level, the individual participant level or both. | |
| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
| Methods – Randomization - implementation | | | |
| 10 Who generated the random allocation sequence, who enrolled participants, and who assigned participants to interventions? | | Replace by 10a, 10b and 10c. 10a Who generated the random allocation sequence, who enrolled clusters, and who assigned clusters to interventions. 10b Mechanism by which individual participants were included in clusters for the purposes of the trial (such as complete enumeration, random sampling). 10c From whom consent was sought (representatives of the cluster, or individual cluster members, or both), and whether consent was sought before or after randomization. | |
| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
| Methods - Blinding | | | |
| 11a If done, who was blinded after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how. 11b If relevant, description of the similarity of interventions. | 11a Whether or not those administering co-interventions were blinded to group assignment. 11b If blinded, method of blinding and description of the similarity of interventions. | | If blinding was not done, or was not possible, explain why. |

From the pool of 1.377 public municipality schools in Denmark, groups of schools are randomly selected by a computer program (STATA) for the trial. The procedure is run by a researcher who works independently of the program developer and intervention staff.

Different implementations strategies are used in the 3 different trial groups (trial groups A-C) as described in the trial design and intervention section above.

There will be no restrictions in the randomization procedure (stratification, matching, blocking).

Statistical Methods

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|--|---|-----------------------|
| Methods - Statistical methods | | | |
| <p>12a Statistical methods used to compare groups for primary and secondary outcomes.</p> <p>12b Methods for additional analyses, such as subgroup analyses and adjusted analyses.</p> | When applicable, details of whether and how the clustering by care providers or centers was addressed. | 12a How clustering was taken into account. | |

Linear and logistic regression models will respectively be used to estimate the impact of the intervention on continuous variables of the primary outcome. The analyses will take into consideration clustering and a few covariates available at baseline, including sex, age, and grade. Intention to treat analyses will be applied to the primary outcomes, as the registers have information on almost all school children

We will do sub-analyses, stratified on sex and grade, since we expect that the impact of interventions will vary by sex and grade.

Time table

| 2013 | 2014 | 2015 | 2016 | 2017 | 2018 |
|---------------|--------------|----------------------------|------|------|---------------------------------------|
| Prepare trial | Intervention | Data collection & analyses | | | Publication & knowledge dissemination |

Ethical and legal consideration

According to Danish Ethical legislation (§ 2 no. 1, law no 593, June 14th, 2011) this trial does not need ethical committee approval.

Applications for trial approval have been send in June 2013 to The Danish Data Protection Agency and the Danish Ministry of Education.

This trial follows the CONSORT – statement criteria including the extensions for Non-pharmacological Treatment Interventions, Cluster Randomized Trials and Pragmatic Trials.

Information to participants in the trial groups: See appendix A and B.

The control group is a “shadow control group” – because the control group schools do not know they are included as control in a scientific trial and because outcome exclusively relies on standard register data collected anyway. The control group continues “Service/Treatment As Usual”. Being in the control group does not restrict their actions or services in any way. Thus the control group schools are not ‘disturbed’ in anyway.

The intervention Robusthed.dk consists of the dissemination of scientifically based knowledge and inspiration for reflection and conversations in the daily setting of family and institutional life. The specific use of the elements in the program is tailored by the responsible adults around the children. No children in the intervention or control groups are prevented from receiving any kind of service as usual as a result of the trials.

Project Feasibility

The intervention program has been specifically designed as a low cost brief intervention program for large scale use. Trial size has been dimensioned to fit with this design and also on the basis of experiences from similar earlier projects (Lundgaard Bak 2012) and pilot experiences.

Qualifications of the research group

The research group has a long experience in primary and public health care work and science –especially within clinical epidemiology and practical method development, evaluation and implementation. For detailed information please refer to the CV’s and the Research Program website www.iupgrowth.com .

Child Mental Health Research Program responsible: Associate professor Phd Carsten Obel

Robusthed.dk trial responsible: Poul Lundgaard Bak, MD

Data-analysis: Jin Liang Zhu, MD

Publication

Results will be published in peer-reviewed journals, at national and international conferences and national media as well as on the research program website. Publication will follow the CONSORT guidelines as described in this protocol. The CONSORT items which pertain to result presentation and discussion in the publications are shown in Appendix C.

Scientific Perspectives

We believe that the intervention project will contribute with knowledge about whether it is possible to support vulnerable children, young people and families on a large scale with low cost programs using contemporary knowledge and technologies.

Practical perspectives, knowledge dissemination

The current program has specifically been designed for easy and low cost implementation. If relevant positive effects can be demonstrated we will recommend starting nationwide implementation via conferences arranged in collaboration with key organizations.

On a national level it is estimated that 1-2 consultants per municipality can implement the program adequately in a community wide setting (public school settings as well as other relevant settings) - on a part time basis integrated as a part of their job portfolio. Using professionals employed in the municipalities as consultants will ensure ownership and anchor the program in the organizations. Based on previous experiences (Lundgaard Bak 2012) it is estimated that 10 day training (including basic course + follow up) is sufficient for a consultant to achieve competency for local implementation. It means that if training courses is held in the 5 regions of the country, implementation can be started on a national scale within ½-1 year after completion of the research project – provided of course that the research results documents that it is worthwhile to implement the program on a larger scale and resources for training of consultants are available.

If the results of the research project show that giving log-in to the program website is “enough” – that is introductory lectures and courses are not necessary, implementation costs are drastically reduced.

In case of sufficient positive effects there will of course also be permanent open access to the website.

Protocol access

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---------------------|-----------------------|
| Other information - protocol | | | |
| 24 Where the full trial protocol can be accessed, if available. | | | |

The full trial protocol will be available (when completed) at The Child Mental Health Research Program website www.iupgrowth.com .

Funding

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---------------------|-----------------------|
| Other information - | | | |
| 25 Sources of funding and other support (such as supply of drugs), role of funders. | | | |

This trial has been funded by The Tryg Foundation.

References

- Adi Y, Killoran A, Janmohamed K, Stewart-Brown S. Systematic review of the effectiveness of interventions to promote mental wellbeing in children in primary education: Report 1: Universal Approaches Non-violence related outcomes. NICE 2007a.
- Adi Y, McMillan AS, Killoran A, Stewart-Brown S. Systematic review of the effectiveness of interventions to promote mental wellbeing in primary schools Report 3: Universal Approaches with focus on prevention of violence and bullying. NICE 2007b.
- Christakis NA, Fowler JH. The Collective Dynamics of Smoking in a Large Social Network. *New England Journal of Medicine* 2008;358(21):2249-58.
- Fowler JH, Christakis NA. Dynamic spread of happiness in a large social network: Longitudinal analysis over 20 years in the Framingham Heart Study. *British Medical Journal* 2008;337:a2338.
- Kadzin AE, Blasé SL. Rebooting Psychotherapy Research and Practice to reduce the Burden of Mental Illness. *Perspectives on Psychological Science* 2011;6(1):21-37.
- Lundgaard Bak, P: Mentalizing communities for children, in Midgley N, Vrouva I (eds.): *Mentalization based interventions with children and families*, Routledge 2012.
- Moffitt TE, Arseneault L, Belsky D, Dickson N, Hancox RJ, Harrington H, Shouts R, Poulton R, Roberts BW, Ross S, Sears MR, Thomson WM, Caspi A. A gradient of childhood self-control predicts health, wealth and public safety. *PNAS* 2011; 108(7): 2693–8.
- Mohr DC, Cuijpers P, Lehman K. Supportive Accountability: A Model for Providing Human Support to Enhance Adherence to eHealth Interventions. *J Med Internet Res* 2011;13(1):e30.
<http://www.jmir.org/2011/1/e30/>
- Montgomery P, Bjornstad G, Dennis J. Media-based behavioral treatments for behavioral problems in children. *Cochrane Database Syst Rev*. 2006 Jan 25;(1):CD002206
- Neil AL, Batterham P, Christensen H, Bennett K, Griffiths KM. Predictors of Adherence by Adolescents to a Cognitive Behavior Therapy Website in School and Community-Based Settings. *J Med Internet Res* 2009;11(1):e6. <http://www.jmir.org/2009/1/e6/>
- NICE: Promoting children’s social and emotional wellbeing in primary education. NICE 2008.
- NICE: Promoting young people’s social and emotional wellbeing in secondary education. NICE 2009.
- O’Connell ME, Boat T, Warner KE. Preventing Mental, Emotional, and Behavioral Disorders among Young People: Progress and Possibilities. National Research Council and Institute of Medicine 2009.
- Ottosen MH, Andersen D, Nielsen LP, Lausten M, Stage S. Børn og Unge i Danmark. Velfærd og Trivsel. SFI 2010.
- Perkins SSJ, Murphy RRM, Schmidt UUS, Williams C. Self-help and guided self-help for eating disorders (Review). *The Cochrane Library* 2009, Issue 1.

Roth A, Fonagy P: What works for whom (2 Ed.). Guildford Press 2006.

Russell E, Koren G, Rieder M, Van Uum S. Hair cortisol as a biological marker of chronic stress: Current status, future directions and unanswered questions. *Psychoneuroendocrinology* 2012; 37:589-601.

Shucksmith J, Summerbell C, Jones S, Whittaker V. Mental wellbeing of children in primary education (targeted/indicated activities). NICE 2007.

The Coalition for Evidence Based Policy: http://coalition4evidence.org/wordpress/?page_id=468
How Low-Cost Randomized Controlled Trials Are Possible in Many Areas of Social Policy:
<http://coalition4evidence.org/wordpress/wp-content/uploads/Rigorous-Program-Evaluations-on-a-Budget-March-2012.pdf>

The Government Office for Science, London: Foresight Mental Capital and Wellbeing Project. 2008.

WHO: What is the evidence on school health promotion in improving health or preventing disease and, specifically, what is the effectiveness of the health promoting schools approach? WHO Europe 2006.

Appendix A: Letter to school leaders in trial group A (in Danish).

Kære skoleleder

Jeg kontakter dig, fordi vi gerne vil tilbyde XX skole adgang til vores videns- og inspirationsprogram Robusthed.dk.

Robusthed handler om at blive god til at klare dagens udfordringer i stort og småt, især når livet er svært. Robusthed kan bruges til lidt af hvert, f.eks.: Gøre det nemmere at lære, træffe gode beslutninger og forebygge stress & konflikter.

Programmet indeholder praktisk viden om robusthed for børn og unge og voksne – i hverdagsprog. Her er både god gammel og ny viden om tanker, følelser og hjernen (også teenage-hjernen), viden om mobning, samt gode historier og små spil. Der er også viden og inspiration i forhold til løsning af en række almindelige og alvorlige typer af problemer. Og man kan bruge programmets smartphone app til praktisk træning.

Robusthed.dk kan betragtes som "videns-vitaminer", som kan flettes ind i små og store situationer i hverdagen, uden at det kræver større indsats end hvad man ellers har mulighed for – måske på et vigtigt tidspunkt i nogle sekunder eller minutter.

Materialet på Robusthed.dk er "grydeklart" og kan fuldstændigt tilpasses det, man i øvrigt arbejder med. Det betyder bl.a., at Robusthed.dk kan integreres i enhver fagpersons samarbejde med et barn, en ung og en familie – idet der er tale om faggruppe- og metodeuafhængig basisviden. En skoleelev, en forælder og en højtuddannet fagperson kan alle have lige stort udbytte. Programmet kan bruges direkte til børn helt ned i 6-7 års alderen. Og forældre og pædagoger kan bruge det i forhold til opdragelse af små børn. Alt på sitet kan både læses og lyttes. Der er også "kursus-forslag" til brug for formaliserede forløb.

Erfaringer viser, at programmet opleves som meningsfuldt af både børn, unge, forældre og professionelle. Oplevelsen er, at det er nemt at bruge, og hjælpsomt i forhold til at løse udfordringer, som fylder i hverdagen. Eksempelvis ser det ud til, at programmet kan bidrage til reduktion af alvorlige konflikter blandt børn og unge med mere end 90 % - og bidrage til betragtelig reduktion i stress-relateret sygefravær hos voksne. Vi mener, at begge dimensioner er relevante i en tid med mange udfordringer.

Robusthed.dk indgår nu i et omfattende forskningsprogram i Danmark og på Grønland.

Vi anbefaler, at log-in deles ud til alle medarbejdere og alle forældre via jeres intranet. De hidtidige erfaringer tyder klart på, at en sådan "lav-intensiv" indsats faktisk er nok til at komme i gang. Forslag til forældre-info er vedhæftet.

Der følger ingen forpligtelser med tilbuddet.

| | | |
|--|----------------|-----------------|
| Log-in til programmet www.robusthed.dk | Brugernavn: XX | Adgangskode: XX |
| Tilhørende I-phone app "Tankespil" downloades fra App store (Android versionen bliver klar i august 2013). | Brugernavn: XX | Adgangskode: XX |

Venlig hilsen

Læge Poul Lundgaard Bak
Forskningsprogrammet for Mental Børnesundhed
Aarhus Universitet og Region Midt

Appendix A.1: Letter to parents from the school leader in participating schools (in Danish).

Kære forældre

Fra Aarhus Universitet har vi fået tilsendt et spændende program, som hedder Robusthed.dk. Alle skolens forældre og alle skolens medarbejdere får fri adgang til programmet.

Robusthed handler om at blive god til at klare dagens udfordringer i stort og småt, især når livet er svært. Robusthed kan bruges til lidt af hvert, f.eks.: Gøre det nemmere at lære, træffe gode beslutninger og forebygge stress & konflikter.

Programmet indeholder praktisk viden om robusthed for børn og unge og voksne – i hverdagsprog. Her er både god gammel og ny viden om tanker, følelser og hjernen (også teenage-hjernen), viden om mobning, samt gode historier og små spil. Der er også viden og inspiration i forhold til løsning af en række almindelige og alvorlige typer af problemer. Det hele kan både læses og lyttes. Og man kan bruge programmets smartphone app til praktisk træning.

Robusthed.dk kan betragtes som "videns-vitaminer", som kan flettes ind i små og store situationer i hverdagen, uden at det kræver større indsats end hvad man ellers har mulighed for – måske på et vigtigt tidspunkt i nogle sekunder eller minutter.

Erfaringer viser, at programmet opleves som meningsfuldt af både børn, unge, forældre og professionelle. Oplevelsen er, at det er nemt at bruge, og hjælpsomt i forhold til at løse udfordringer, som fylder i hverdagen. Eksempelvis ser det ud til, at programmet kan bidrage til reduktion af alvorlige konflikter blandt børn og unge med mere end 90 % - og bidrage til betragtelig stress reduktion hos voksne. Vi mener, at begge dimensioner er relevante i en tid med mange udfordringer.

Robusthed.dk indgår nu i et omfattende forskningsprogram i Danmark og på Grønland.

Her er adgangsoplysninger til programmet:

| | | |
|--|----------------|-----------------|
| Log-in til programmet www.robusthed.dk | Brugernavn: XX | Adgangskode: XX |
| Tilhørende I-phone app "Tankespil" downloades fra App store (Android versionen bliver klar i august 2013). | Brugernavn: XX | Adgangskode: XX |

Venlig hilsen

XX

Skoleleder

Appendix B: Letter to school leaders in trial group B (in Danish).

Kære skoleleder

Jeg kontakter dig, fordi vi gerne vil tilbyde XX skole adgang til vores videns- og inspirationsprogram Robusthed.dk.

Robusthed handler om at blive god til at klare dagens udfordringer i stort og småt, især når livet er svært. Robusthed kan bruges til lidt af hvert, f.eks.: Gøre det nemmere at lære, træffe gode beslutninger og forebygge stress & konflikter.

Programmet indeholder praktisk viden om robusthed for børn og unge og voksne – i hverdagsprog. Her er både god gammel og ny viden om tanker, følelser og hjernen (også teenage-hjernen), viden om mobning, samt gode historier og små spil. Der er også viden og inspiration i forhold til løsning af en række almindelige og alvorlige typer af problemer. Og man kan bruge programmets smartphone app til praktisk træning.

Robusthed.dk kan betragtes som "videns-vitaminer", som kan flettes ind i små og store situationer i hverdagen, uden at det kræver større indsats end hvad man ellers har mulighed for – måske på et vigtigt tidspunkt i nogle sekunder eller minutter.

Materialet på Robusthed.dk er "grydeklart" og kan fuldstændigt tilpasses det, man i øvrigt arbejder med. Det betyder bl.a., at Robusthed.dk kan integreres i enhver fagpersons samarbejde med et barn, en ung og en familie – idet der er tale om faggruppe- og metodeuafhængig basisviden. En skoleelev, en forælder og en højtuddannet fagperson kan alle have lige stort udbytte. Programmet kan bruges direkte til børn helt ned i 6-7 års alderen. Og forældre og pædagoger kan bruge det i forhold til opdragelse af små børn. Alt på sitet kan både læses og lyttes. Der er også "kursus-forslag" til brug for formaliserede forløb.

Erfaringer viser, at programmet opleves som meningsfuldt af både børn, unge, forældre og professionelle. Oplevelsen er, at det er nemt at bruge, og hjælpsomt i forhold til at løse udfordringer, som fylder i hverdagen. Eksempelvis ser det ud til, at programmet kan bidrage til reduktion af alvorlige konflikter blandt børn og unge med mere end 90 % - og bidrage til betragtelig reduktion i stress-relateret sygefravær hos voksne. Vi mener, at begge dimensioner er relevante i en tid med mange udfordringer.

Robusthed.dk indgår nu i et omfattende forskningsprogram i Danmark og på Grønland.

Vi vil gerne tilbyde intro-foredrag i det omfang, som måtte passe jer, samt log-in til Robusthed.dk. Vi anbefaler, at log-in deles ud til alle medarbejdere og alle forældre via jeres intranet. De hidtidige erfaringer tyder klart på, at en sådan "lav-intensiv" indsats faktisk er nok til at komme i gang. Forslag til forældre-info er vedhæftet.

Der følger ingen forpligtelser med tilbuddet – men naturligvis et tilbud om opfølgning, hvis I ønsker det.

| | | |
|--|-------------------|--------------------|
| Log-in til programmet www.robusthed.dk | Brugernavn: nydam | Adgangskode: nydam |
| Tilhørende I-phone app "Tankespil" downloades fra App store (Android versionen bliver klar i august 2013). | Brugernavn: nydam | Adgangskode: nydam |

Jeg tillader mig at kontakte dig pr telefon indenfor den kommende måned for at drøfte mulighederne for samarbejde, som beskrevet ovenfor. Du er også velkommen til at kontakte mig på mail eller tlf. 31 26 06 78.

Læge Poul Lundgaard Bak
Forskningsprogrammet for Mental Børnesundhed
Aarhus Universitet og Region Midt

Appendix C: CONSORT items which will be followed in the presentation of results and discussion in publications of the trial:

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---|--|
| Results – participant flow (download diagram) | | | |
| 13a For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analyzed for the primary outcome. 13b For each group, losses and exclusions after randomization, together with reasons. | The number of care providers or centers performing the intervention in each group and the number of patients treated by each care provider or in each center. | 13a For each group, the numbers of clusters that were randomly assigned, received intended treatment, and were analyzed for the primary outcome. 13b For each group, losses and exclusions for both clusters and individual cluster members. | The number of participants or units approached to take part in the trial, the number which was eligible, and reasons for non-participation should be reported. |
| Implementation of intervention | | | |
| | Details of the experimental treatment and comparator as they were implemented. | | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|---|---------------------|-----------------------|
| Results – Recruitment | | | |
| 14a Dates defining the periods of recruitment and follow-up. 14b Why the trial ended or was stopped. | | | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|--|--|-----------------------|
| Results – baseline data | | | |
| 15 A table showing baseline demographic and clinical characteristics for each group | When applicable, a description of care providers (case volume, qualification, expertise, etc.) and centers (volume) in each group. | Baseline characteristics for the individual and cluster levels as applicable for each group. | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---|-----------------------|
| Results – numbers analyzed | | | |
| 16 For each group, number of participants (denominator) included in each analysis and whether the analysis was by original assigned groups. | | For each group, number of clusters included in each analysis. | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---|-----------------------|
| Results – Outcomes and estimation | | | |
| 17a For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval). 17b For binary outcomes, presentation of both absolute and relative effect sizes is recommended. | | 17a Results at the individual or cluster level as applicable and a coefficient of intra-cluster correlation (ICC or k) for each primary outcome. | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---------------------|-----------------------|
| Results – ancillary analyses | | | |
| 18 Results of any other analyses performed, including subgroup analyses and adjusted analyses, distinguishing pre-specified from exploratory. | | | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---------------------|-----------------------|
| Results – Harms | | | |
| 19 All important harms or unintended effects in each group (for specific guidance see CONSORT for harms). | | | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|---|---------------------|-----------------------|
| Discussion – limitations | | | |
| 20 Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses. | | | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|--|--|---|
| Discussion – generalizability | | | |
| 21 Generalizability (external validity, applicability) of the trial findings. | Generalizability (external validity) of the trial findings according to the intervention, comparators, patients, and care providers and centers involved in the trial. | Generalizability to clusters and/or individual participants (as relevant). | Describe key aspects of the setting which determined the trial results. Discuss possible differences in other settings where clinical traditions, health service organization, staffing, or resources may vary from those of the trial. |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|--|---|---------------------|-----------------------|
| Discussion – Interpretation | | | |
| 22 Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence. | 20 In addition, take into account the choice of the comparator, lack of or partial blinding, and unequal expertise of care providers or centers in each group. | | |

| CONSORT – Statement (CON) | Non-Pharmacological Treatment Interventions (NPT) | Cluster Trials (CT) | Pragmatic Trials (PT) |
|---|---|---------------------|-----------------------|
| Other information - registration | | | |
| 23 Registration number and name of trial registry. | | | |